



# The Dyskinetic Cerebral Palsy Functional Impact Scale: development and validation of a new tool

KIRSTY STEWART<sup>1,2,3</sup>  | JENNIFER LEWIS<sup>1</sup> | MARGARET WALLEN<sup>3</sup>  | NATASHA BEAR<sup>4</sup> | ADRIENNE HARVEY<sup>5</sup>

**1** Kids Rehab, The Children's Hospital at Westmead, Sydney, New South Wales; **2** Faculty of Medicine and Health, The University of Sydney, Sydney, New South Wales; **3** School of Allied Health, Australian Catholic University, North Sydney, New South Wales; **4** Institute of Health Research, University of Notre Dame, Fremantle, Western Australia; **5** Neurodisability and Rehabilitation, Murdoch Children's Research Institute, Melbourne, Victoria, Australia.

Correspondence to Kirsty Stewart at Kids Rehab, The Children's Hospital at Westmead, Locked bag 4001, Westmead, NSW 2145, Australia. E-mail: kirsty.stewart@health.nsw.gov.au

This original article is commented by Himmelmann on page 1370 of this issue.

## PUBLICATION DATA

Accepted for publication 19th May 2021.

Published online 19th June 2021.

## ABBREVIATIONS

BADS	Barry-Albright Dystonia Scale
CFCS	Communication Function Classification System
D-FIS	Dyskinetic Cerebral Palsy Functional Impact Scale
EDACS	Eating and Drinking Ability Classification System
GMFCS	Gross Motor Function Classification System
MACS	Manual Ability Classification System

**AIM** To outline the development and examine the content and construct validity of a new tool, the Dyskinetic Cerebral Palsy Functional Impact Scale (D-FIS), which measures the impact of dyskinesia on everyday activities in children with cerebral palsy (CP).

**METHOD** D-FIS content was informed by a systematic review of dyskinesia outcome measures, in collaboration with children with dyskinetic CP, parents, caregivers, and expert clinicians. The D-FIS uses parent proxy to rate impact of dyskinesia on everyday activities. Construct validity was determined by examining internal consistency; known groups validity with the Gross Motor Function Classification System (GMFCS), Manual Ability Classification System (MACS), Communication Function Classification System (CFCS), and Eating and Drinking Ability Classification System (EDACS); and convergent validity with the Barry-Albright Dystonia Scale (BADS).

**RESULTS** Fifty-seven parents of children (29 males, 28 females, mean [SD] age 11y 8mo [4y 4mo], range 2y 6mo–18y) completed the D-FIS. Correlation between D-FIS and GMFCS was  $r=0.86$  (95% confidence interval [CI]: 0.77–0.91,  $p<0.001$ ); MACS  $r=0.84$  (95% CI: 0.73–0.90,  $p<0.001$ ); CFCS  $r=0.80$  (95% CI: 0.67–0.88,  $p<0.001$ ); and EDACS  $r=0.78$  (95% CI: 0.66–0.87). Correlation between D-FIS and BADS was  $r=0.77$  (95% CI: 0.64–0.86,  $p<0.001$ ). Cronbach's alpha was 0.96.

**INTERPRETATION** The D-FIS demonstrates good construct validity and high internal consistency. The D-FIS will be useful for identifying priorities for intervention. It adds to the measurement tool kit for children with dyskinetic CP by addressing functional impact of dyskinetic movements and postures.

Dyskinetic cerebral palsy (CP), one of the most disabling forms of CP,<sup>1,2</sup> is a motor disorder characterized by changes in muscle tone and posture, with varying degrees of involuntary movement.<sup>3</sup> Dystonia and choreoathetosis are the two clinical manifestations of dyskinetic CP, although many children classified as dyskinetic CP also present with spasticity.<sup>4,5</sup> Standardized and accurate measurement of dyskinesia in people with CP is important to objectively quantify the movement disorder, monitor intervention outcomes, ensure practice is based on high quality evidence, and guide future interventions.

A number of scales have been developed to measure dystonia and choreoathetosis severity in children with dyskinetic CP.<sup>6</sup> The majority of scales assess dystonia severity only, taking into consideration its duration, provoking factor, and amplitude. The most recently published Dyskinesia Impairment Scale<sup>7</sup> is the only scale to address both dystonia and choreoathetosis in CP and the Barry-Albright Dystonia Scale (BADS)<sup>8</sup> is the only other scale developed

specifically for people with CP. All dyskinesia scales are intended as impairment, body function, and structure level assessments according to the International Classification of Functioning, Disability and Health (ICF),<sup>9</sup> with some providing limited insight into the impact of the dyskinesia or movement disorder on a small number of broad daily activities. In addition to severity, the importance of assessing the activity and participation domains of the ICF, considering individual personal, cultural, and environmental factors, has been highlighted.<sup>10–12</sup>

Dyskinesia scales for people with CP can be used for two purposes: to provide a measure of severity at a single time point and to measure change after an intervention targeting dyskinesia.<sup>6</sup> However, a reduction in dyskinesia severity does not necessarily translate to enhanced function or participation, or improvements in caregiver assistance and quality of life.<sup>13</sup> A clinical tool which will assist in identifying how changes in dyskinesia severity translates to changes in function, activity participation, caregiver

assistance, and thus impact quality of life would be a useful adjunct to clinical and research practice.

A number of well-established, psychometrically robust outcome measures currently exist for use with children with CP and with disability generally.<sup>14–21</sup> When examining baseline function and measuring intervention outcomes of children with dyskinetic CP, multiple questionnaires and outcome tools are required to adequately cover all domains of the ICF,<sup>22</sup> few of which have been developed specifically for children with dyskinetic CP and the unique functional limitations they may experience. This may result in assessment burden and fatigue for families, children, and clinicians. In addition, children with dyskinesia are typically seen in busy clinics where assessment needs to be feasible, focused on functional activities, and enables family priorities for intervention to be identified and measured efficiently and effectively. A single tool to specifically measure the impact of dyskinesia on daily function in children with dyskinetic CP is therefore considered a valuable addition to the existing measures.

The Dyskinetic Cerebral Palsy Functional Impact Scale (D-FIS) was developed to identify the impact of dyskinesia on daily functional activities for children and their caregivers and guide families in identifying daily activities which are their priorities for intervention.

The aim of this paper is to describe the development of the D-FIS and to report on a study evaluating its content and construct validity.

## METHOD

### Development of the D-FIS

The need for a new tool to specifically quantify the impact of dyskinesia on daily function in children with CP was identified: (1) by parents of children with dyskinetic CP and children with dyskinetic CP during a qualitative study investigating the lived experience of dyskinetic CP;<sup>23</sup> and (2) during an Australian Dyskinetic CP symposium in 2015 attended by occupational therapists, physiotherapists, speech pathologists, paediatric rehabilitation physicians, and paediatric neurologists ( $n=115$ ).

Development of the D-FIS as an evaluative tool followed the methodological framework for assessing health indices by Kirshner and Guyatt.<sup>24</sup> The items for the pilot version, aimed at comprehensively covering usual daily activities for children, were generated from: (1) the extensive clinical experience of the research team; (2) recommendations obtained by interview, from parents and caregivers of children with dyskinetic CP, and children and adolescents with dyskinetic CP; and (3) a survey of 115 clinicians with interest and expertise in the management of children with dyskinetic CP attending the National Dyskinesia Symposium.<sup>23</sup> Daily activities and constructs deemed most important to families, children, and clinicians, and that were part of daily routine, formed the item pool. The item pool was further informed by current best evidence in the assessment of dyskinesia,<sup>23</sup> the results of a systematic review of currently available dyskinesia measures,<sup>6</sup> and literature in

### What this paper adds

- The Dyskinetic Cerebral Palsy Functional Impact Scale (D-FIS) assesses the perceived impact of dyskinesia on daily activities in children with cerebral palsy (CP).
- The D-FIS demonstrates good construct validity and high internal consistency.
- The D-FIS is a clinically feasible, family-centred tool that fills a current gap in the dyskinetic CP assessment toolkit.

the areas of quality of life and caregiver burden,<sup>15,20</sup> family-centred goal setting,<sup>14</sup> sleep,<sup>18</sup> pain,<sup>16,21</sup> and participation.<sup>17,25</sup>

A 16-item pilot version was subjected to review and feedback to develop and refine the D-FIS. A convenience sample of eight primary caregivers (six females, two males) of children with dyskinetic CP, recruited from an outpatient clinic of the Children's Hospital at Westmead, completed the pilot D-FIS. These primary caregivers reported for six children (mean age 14y 1mo [range 5y 7mo–18y 6mo, SD 4y 10mo], Gross Motor Function Classification System [GMFCS]<sup>26</sup> levels: II=1, III=1, IV=3, V=1; Manual Ability Classification System [MACS]<sup>27</sup> levels: II=1, III=2, IV=2, V=1; Communication Function Classification System [CFCS]<sup>28</sup> levels: I=3, III=1, IV=2; and the Eating and Drinking Ability Classification System [EDACS] levels: I=3, III=1, V=2).<sup>29</sup> This convenience sample broadly represents CP classification levels and the general age range of children with dyskinetic CP for whom this tool is intended. Parents then participated in semi-structured face-to-face interviews to elicit views on the D-FIS and identify revisions. The pilot version of the D-FIS was also emailed to a purposive sample of eight Australian clinicians with specific expertise in the management of children with dyskinetic CP and/or experience in outcome measurement development to elicit written feedback on the items, wording, scales, and coverage of the D-FIS. Clinicians included paediatric rehabilitation physicians, a CP sleep disturbance expert, occupational, physical, and speech therapists, and a paediatric neurologist. These combined processes resulted in two additional items, inclusion of the Priority Scale, refined content and language of all items and scales, and enhanced usefulness of the questionnaire. The final 18-item D-FIS is under exploration in this study.

All primary caregiver participants recruited for the study were provided with the printed questionnaire and detailed instructions. The front page of the D-FIS explains dyskinesia as it is seen in children with CP. The questionnaire was accompanied, in all cases, by a discussion regarding: (1) their child's specific movement disorders; (2) how to identify dyskinetic movements and postures in their child; and (3) that dyskinetic movements and postures may be only one reason why their child may have difficulty with daily activities. We also reminded families that there may be alternative reasons for a functional difficulty – each item has a 'not applicable' option with the following wording: 'Standing (depending on the item) is difficult but not due to dyskinesia'.

Development of the D-FIS progressed in stages and was approved by the Royal Children's Hospital (RCH HREC: 36129A) and the Sydney Children's Hospitals Network (LNR/18/SCHN/32) Human Research Ethics Committees.

### The D-FIS

The final D-FIS consists of 18 items: 16 daily activities and two additional constructs that impact children with dyskinetic CP (Pain and Fatigue). It takes approximately 10 minutes to complete via parent proxy report (Appendix S1, online supporting information). The D-FIS items are rated on two scales: an Impact Scale and a Priority Scale. Each item is a separate functional activity: Sitting; Standing; Walking; Positioning; Transfers; Other Gross Motor Activities; Fine Motor Activities; Reaching; Use of Technology; Daily Hygiene Activities; Upper Body Dressing; Lower Body Dressing; Feeding; Speech; Sleep; and Leisure Activities. Pain and Fatigue make up the two additional constructs. The Impact Scale rates the impact dyskinesia has on each activity/construct on a 5-point ordinal scale from 0 (no impact), 1 (mild impact), 2 (moderate impact), 3 (severe impact), to 4 (extreme impact). The Impact Scale ratings also account for functioning in each activity without the assistance of equipment and/or a caregiver and function with the assistance of equipment and/or caregiver. Each activity is briefly described in the tool and examples are provided that span developmental levels. These examples are not intended to be all-inclusive but rather to give parents an understanding of what the activity may cover. The Impact Scale scores are summed, the total score ranges from 0 to 72, and higher scores indicate greater impact of dyskinesia on daily functional activities. The Priority Scale identifies the current priority of each activity for the child and their family from 1 (not a priority) to 4 (highest priority). The purpose of the Priority Scale is to assist families and clinicians with goal identification and ensuring the activity areas most important to children and their families are the focus of intervention.

### Construct validity of the D-FIS

#### Participants

A convenience sample of primary caregivers ( $n=57$ ) were recruited from outpatients attending the Kids Rehab Department of the Children's Hospital at Westmead between October 2018 and November 2019. Eligible caregivers were those with children aged 2 to 18 years with dyskinetic CP, or dystonia/choreoathetosis and spasticity where dystonia or choreoathetosis was the predominant motor type. Predominance of motor type was determined clinically using the Cerebral Palsy Description Form.<sup>30</sup> Forty-nine (86%) of the 57 primary caregivers were mothers or female carers. Their children were aged between 2 years 6 months and 18 years (mean age 11y 7mo, SD 4y 4mo). Functional classification levels (GMFCS, MACS, CFCS, and EDACS) were assigned for all participants and

comorbidities recorded (Table 1). Informed, written consent was obtained from all participating caregivers.

Construct validity, the degree to which the D-FIS measures the intended construct, was evaluated using known groups methods based on correlation of the D-FIS with the functional levels of the GMFCS, MACS, CFCS, and EDACS, and differences in D-FIS scores across levels within each of these four systems. Each system classifies children across five levels, with level I indicating minimal disability and a high level of independence and level V indicating total dependence on equipment and carers for all daily needs. Differences in D-FIS total scores between ambulant and non-ambulant children according to the GMFCS (ambulant=levels I, II, and III; non-ambulant=levels IV and V) were compared. The hypotheses were that: (1) children in higher GMFCS/MACS/

**Table 1:** Participant demographics and classification levels of their child with cerebral palsy (CP)

Demographics	
Age (y:mo)	
Range	2:6–18:0
Mean (SD)	11:7 (4:4)
Sex	
Male	29 (51)
Female	28 (49)
Distribution	
Bilateral CP	46 (80)
Unilateral CP	11 (20)
GMFCS level	
I	8 (14)
II	6 (11)
III	7 (12)
IV	23 (40)
V	13 (23)
MACS level	
I	0
II	19 (33)
III	12 (21)
IV	12 (21)
V	14 (25)
CFCS level	
I	16 (28)
II	13 (23)
III	14 (25)
IV	10 (17)
V	4 (7)
EDACS level	
I	16 (28)
II	19 (33)
III	11 (19)
IV	1 (2)
V	10 (17)
Comorbidities	
Nil	13 (23)
Epilepsy	16 (28)
Gastrostomy	9 (16)
Visual difficulties	16 (28)
Hearing impairment	4 (7)
Intellectual disability	23 (40)

Data are  $n$  (%) unless otherwise stated. GMFCS, Gross Motor Function Classification System; MACS, Manual Ability Classification System; CFCS, Communication Function Classification System; EDACS, Eating and Drinking Ability Classification System.

CFCS/EDACS levels (children with lower function) would have higher D-FIS Impact Scale total scores (lower ability in daily functional activities); (2) D-FIS would be strongly positively correlated with the GMFCS and MACS; (3) D-FIS would be moderately correlated with the CFCS and EDACS; and (4) ambulant children would have significantly lower D-FIS scores than non-ambulant children.

Construct convergent validity,<sup>31</sup> the extent to which the D-FIS produces similar results as another well-established tool that measures a related construct (i.e. dystonia severity), was assessed against the BADS. The BADS was selected as it is a clinically useful scale, developed specifically for people with CP, has demonstrated responsiveness to change after interventions, and is used extensively in Australia.<sup>32</sup> The BADS takes approximately 10 minutes to complete which minimizes child and family assessment burden. The BADS scores dystonia on a 5-point, criterion-based, ordinal severity scale (0–4) across eight body regions. Scores for each body region are summed to a maximum of 32, with higher scores indicating increasing severity of dystonia. The BADS was completed with every child by two of the researchers (KS, JL) who have extensive experience rating dyskinesia in children with CP. The hypothesis was that children with higher BADS scores would have higher D-FIS Impact Scale total scores, and these would be strongly positively correlated.

### Data analysis

Spearman's rank correlation assessed the correlation between total D-FIS Impact Scale scores and the GMFCS, MACS, CFCS, EDACS, and BADS. Confidence intervals (CIs) for Spearman's rank correlation were calculated, based on Fisher's transformation. Correlation coefficient of  $r \geq 0.7$  was considered strong and 0.5 to  $<0.7$  was considered moderate.<sup>33</sup> One-way analysis of variance was used to analyse differences in total D-FIS scores across levels of the classification systems and the difference in D-FIS scores between ambulant and non-ambulant was examined using an independent *t*-test. The D-FIS total score was normally distributed as shown by histograms and box plots and confirmed by the Shapiro–Wilk test and a normal Q–Q plot. Homogeneity of the variances was demonstrated

using Levene's test. A two-tailed significance level of 0.05 was used in all tests. All data were analysed using Stata version 16.1 (StataCorp, College Station, TX, USA).

Ratings on the Priority Scale, intended to focus clinical intervention on those activities rated most important to families, were reported using descriptive statistics. Priority Scale mean scores are presented for ambulant and non-ambulant children.

### RESULTS

The D-FIS was completed by 57 caregivers for 57 children (see Table 1 for characteristics of the children). Children were distributed across the functional classification levels except MACS level I – a finding similar to other dyskinetic CP cohort studies.<sup>5,34</sup>


Mean total D-FIS Impact Scale scores by classification levels are reported in Table 2. The mean total score for the whole cohort ( $n=57$ ) on the D-FIS Impact Scale was 40.44 (SD 16.72, range 8–72). Increasing functional severity levels on the GMFCS, MACS, CFCS, and EDACS demonstrated increased mean D-FIS scores ( $p<0.001$ ). Spearman's rank correlation for D-FIS Impact Scale total score and GMFCS was  $r=0.86$  (95% confidence interval [CI]: 0.77–0.91,  $p<0.001$ ); MACS  $r=0.84$  (95% CI: 0.73–0.90,  $p<0.001$ ); CFCS  $r=0.80$  (95% CI: 0.67–0.88,  $p<0.001$ ); and EDACS  $r=0.78$  (95% CI: 0.66–0.87,  $p<0.001$ ), indicating strong relationships with the D-FIS. Mean total D-FIS Impact Scale scores between ambulant and non-ambulant children were significantly different (respectively  $n=22$ , 23.6 [SD 9.8];  $n=35$ , 51.0 [SD 10.1];  $p<0.001$ ). The correlation between D-FIS Impact Scale total scores and BADS was  $r=0.77$  (95% CI: 0.64–0.86,  $p<0.001$ ), indicating a strong relationship. The D-FIS demonstrated good internal consistency ( $\alpha=0.96$ ).

The mean Priority Scale scores for the ambulant, non-ambulant, and total cohorts are presented in Figure 1. In the ambulant cohort, the highest priority activity, identified by parents for their children, was fine motor activities (mean 3.5) followed by leisure, daily hygiene, gross motor, and walking (means 2.7–3). The least prioritized activities were sleep, sitting, and positioning (total means of 1.6–1.8). In the non-ambulant cohort, the activities rated the

**Table 2:** Mean total scores for the D-FIS Impact Scale by classification levels

Level <sup>a</sup>	Total D-FIS Impact Scale scores <sup>b</sup> for each classification system			
	GMFCS	MACS	CFCS	EDACS
I	17.2 (6.2, 8–27)	0	22.4 (10.8, 8–43)	23.1 (11.2, 8–46)
II	22.4 (7.7, 11–39)	22.5 (9.9, 8–47)	36.5 (10.1, 20–50)	38.8 (11.4, 20–61)
III	32.0 (9.8, 19–49)	38.3 (6.8, 31–47)	47.3 (9.5, 30–61)	49.7 (8.3, 30–58)
IV	46.6 (7.4, 31–61)	52.9 (8.3, 38–69)	59.1 (8.6, 44–72)	$n=1$ child, 69
V	58.5 (9.7, 38–72)	55.9 (9.7, 38–72)	54.5 (12.9, 38–69)	58.3 (10, 38–72)
ANOVA	$F(4,52)=45.4$ $p<0.001$	$F(3,53)=46.6$ $p<0.001$	$F(4,52)=25.1$ $p<0.001$	$F(4,52)=22.0$ $p<0.001$

Data are mean (SD, range) unless otherwise stated. <sup>a</sup>Level I represents higher function, level V represents lower function. <sup>b</sup>Higher D-FIS scores indicate lower function. GMFCS, Gross Motor Function Classification System; MACS, Manual Ability Classification System; CFCS, Communication Function Classification System; EDACS, Eating and Drinking Ability Classification System; ANOVA, analysis of variance.



**Figure 1:** Parent report activity priority for ambulant, non-ambulant, and entire cohort. GMFCS, Gross Motor Function Classification System.

highest priority were transfers (mean 3.65), positioning, technology use, dressing, and daily hygiene activities (means 3.3–3.6). The least important activity rated by parents was gross motor activities (mean 2.3), walking, speech, and fatigue (total means of 2.5–2.8).

## DISCUSSION

This study reports the development and initial psychometric properties of the D-FIS, a newly developed tool to identify the impact of dyskinesia on daily activities in children with CP. The D-FIS was intended to be clinically feasible to implement within the confines of specialty CP clinics in busy children’s and rehabilitation services. The D-FIS demonstrates good content and construct validity and strong internal consistency in children aged 2 years 6 months to 18 years. Dyskinesia is a complex and disabling movement disorder in people with CP. Tools to measure the severity of the movement disorder have been available for decades, yet no single tool assesses the impact of dyskinesia on the daily functional activities of children with CP nor measures the impact of targeted interventions on those functional tasks, either from a parent/caregiver’s or child’s perspective. The D-FIS aims to fill this current gap in measurement.

The D-FIS consists of 16 daily functional activities and two additional constructs, Pain and Fatigue. These additional constructs, although not functional activities, are pain and fatigue are impacted by dyskinesia<sup>11,21,35</sup> and have

been demonstrated to impact a child’s participation across all daily activities. Despite appearing as an additional construct, results indicate they did not impact the internal consistency of the D-FIS.

The D-FIS is able to distinguish between ambulant and non-ambulant children, with a clear gradient of mean total scores across GMFCS and MACS levels. These findings confirm our hypotheses and support the construct validity of the D-FIS. The findings also support the validity of the D-FIS in a heterogenous sample of children in relation to age and severity of presentation. Although analysis of variance identified differences between levels for the CFCS and EDACS, small participant numbers in some levels on these classifications mitigated against observing a gradient of increasing scores. Considered together with the strong correlations between D-FIS total scores and each of the functional classification systems, there is good evidence of construct validity. This supports our initial hypothesis that children with CP with more severe dyskinesia demonstrate poorer function across the activities they complete on a daily basis and require greater physical support from caregivers.

The Priority Scale results demonstrate that activities that require the most physical assistance during care, such as positioning, daily hygiene, and dressing, were amongst the highest priority for the parents of non-ambulant children. Access to technology was also rated as a high priority for this group of children, reflecting the possible reliance on

technology for both school-based academic tasks, communication, and as a quiet time leisure activity. Parents of ambulant children indicated that fine motor activities were the highest priority for their children, which is consistent with the knowledge that dyskinesia frequently affects the upper limbs more than the lower limbs in more ambulant children with CP and in particular children with unilateral involvement.<sup>5</sup> We expected that pain, sleep, and fatigue may feature more frequently as a priority for families, given the increasing focus on these factors in the literature and their impact on quality of life.<sup>11</sup> As the D-FIS is a parent proxy questionnaire, the priority scale reflects the parent's perceptions of their child's experiences, particularly in regards to pain, fatigue, and to some extent sleep, and may not be viewed as a higher priority unless impacting family life. As expected, however, they were reported more frequently as a priority in non-ambulant children. Children themselves may rate these factors as a high priority and these factors may become more prominent with increasing age.

There are factors, other than dyskinesia, which may impact the domains of the D-FIS. Throughout the process of developing the tool, parents were found to be very good at understanding their child's difficulties and the reasons for these difficulties. For example a child may have difficulty accessing their eye gaze system due to their low tone making head control difficult and their intellectual disability and not necessarily their abnormal movements. We ask respondents for their perception of functional ability attributable to dyskinesia. We do not ask families to distinguish between dystonia or choreoathetosis. We have not detected any uncertainty by families in using the individual scales within the D-FIS, which have now been completed as part of the development and validation of the tool. Ultimately, this is a parent proxy tool, which reflects parent perception of their child's ability framed within the context provided by the D-FIS.

The D-FIS authors have found the questionnaire to be responsive and clinically useful in children aged over 3 years. By the age of 3 years, typically developing children are independent in most D-FIS daily functional activities with emerging independent dressing skills and daily hygiene skills, ensuring that the descriptive scale levels for each activity rate the impact of dyskinesia on that activity rather than risk rating developmental ability.

## REFERENCES

- Himmelman K, Hagberg G, Wiklund LM, Eek MN, Uvebrant P. Dyskinetic cerebral palsy: a population based study of children born between 1991 and 1998. *Dev Med Child Neurol* 2007; **49**: 246–51.
- Pr el M, Rackauskaite G, Larsen ML, et al. Children with dyskinetic cerebral palsy are severely affected as compared to bilateral spastic cerebral palsy. *Acta Paediatr* 2019; **108**: 1850–6.
- Platt MJ, Krageloh-Mann I, Cans C. Surveillance of cerebral palsy in Europe: reference and training manual. *Medical Edu* 2009; **43**: 495–6.
- Himmelman K, McManus V, Hagberg G, Uvebrant P, Kr ageloh-Mann I, Cans C. Dyskinetic cerebral palsy in Europe: trends in prevalence and severity. *Arch Dis Child* 2009; **94**: 921–6.
- Sun D, Wang Q, Hou M, et al. Clinical characteristics and functional status of children with different subtypes of dyskinetic cerebral palsy. *Medicine* 2018; **97**: e10817.
- Stewart K, Harvey A, Johnston LM. A systematic review of scales to measure dystonia and choreoathetosis in children with dyskinetic cerebral palsy. *Dev Med Child Neurol* 2017; **59**: 786–95.
- Monbaliu E, Ortibus ELS, de Cat J, et al. The Dyskinesia Impairment Scale: a new instrument to measure dystonia and choreoathetosis in dyskinetic

This study was limited by sample size as recruitment was reliant on clinical availability of families with children with dyskinetic CP attending outpatient clinics. Test–retest reliability is ongoing and, whilst important, is more difficult to determine in a condition such as dyskinetic CP because of the changing nature of the movement disorder and the daily influences of health, environment, mood, and sleep on dyskinesia. At this stage, the D-FIS remains a parent proxy questionnaire. A self-report version is under development.

The D-FIS is valuable as a valid and time-efficient means to identify the impact of dyskinesia on function and to understand family priorities for intervention. Further research is underway to evaluate its responsiveness to change after intrathecal baclofen therapy, deep brain stimulation therapy, and medication trials.

The D-FIS is a newly developed tool that measures the impact of dyskinesia on typical daily functional activities of children with dyskinetic CP. This parent proxy version demonstrates strong validity. Further research is ongoing to determine test–retest reliability and responsiveness to change which will inform use of the D-FIS as an outcome measure. A self-report version will be developed with guidance from children with dyskinetic CP and will provide a more complete picture of the child's function and priorities.

## ACKNOWLEDGEMENTS

The authors wish to acknowledge the valuable contribution to the following parents of children with dyskinetic CP, adolescents with dyskinetic CP, and clinical experts in the field of CP management: J Kelderman, J Kelly, D Kelly, R Shah, A Shah, S Wick, J Charlesworth, M Jordon, S Clifton-Bligh, L Conn, M Conn, Dr S McIntyre, Dr L Copeland, Dr M-C Waugh, N Smith, Dr S McCabe, C Brady, Dr J Rice, and Dr S Mohammad. We also want to acknowledge the Australian National Health and Medical Research Council-funded Centre of Research Excellence in Cerebral Palsy (#1057997). The authors declare no conflicts of interest with respect to the research, authorship, or publication of this article.

## SUPPORTING INFORMATION

The following additional material may be found online:

**Appendix S1:** Dyskinetic Cerebral Palsy Functional Impact Scale.

- cerebral palsy. *Dev Med Child Neurol* 2012; **54**: 278–83.
8. Barry MJ, VanSwearingen JM, Albright AL. Reliability and responsiveness of the Barry-Albright dystonia scale. *Dev Med Child Neurol* 1999; **41**: 404–11.
  9. World Health Organization. International Classification of Functioning, Disability and Health: ICF. Geneva: World Health Organization, 2001.
  10. Harvey A. Challenges and advancements in measuring dyskinesia in cerebral palsy. *Dev Med Child Neurol* 2020; **62**: 411.
  11. Lumsden DE, Gimeno H, Tustin K, Kaminska M, Lin J-P. Interventional studies in childhood dystonia do not address the concerns of children and their carers. *Eur J Paed Neurol* 2015; **19**: 327–36.
  12. Himmelmann K. The quest for patterns in dyskinetic cerebral palsy. *Dev Med Child Neurol* 2016; **58**: 112.
  13. Elkaim LM, De Vloop P, Kalia SK, Lozano AM, Ibrahim GM. Deep brain stimulation for childhood dystonia: current evidence and emerging practice. *Expert Rev Neurother* 2018; **18**: 773–84.
  14. Carlberg EB, Lowing K. Does goal setting in activity-focused interventions for children with cerebral palsy influence treatment outcomes? *Dev Med Child Neurol* 2013; **55**: 47–54.
  15. Carlon S, Shields N, Yong K, Gilmore R, Sakzewski L, Boyd R. A systematic review of the psychometric properties of Quality of Life measures for school aged children with cerebral palsy. *BMC Ped* 2010; **10**: 81.
  16. Doralp S, Bartlett DJ. The prevalence, distribution, and effect of pain among adolescents with cerebral palsy. *Pediatr Phys Ther* 2010; **22**: 26–33.
  17. Imms C. Children with cerebral palsy participate: a review of the literature. *Dis Rehabil* 2008; **30**: 1867–84.
  18. McCabe SM, Blackmore AM, Abbiss CR, Langdon K, Elliott C. Sleep concerns in children and young people with cerebral palsy in their home setting. *J Paed Child Health* 2015; **51**: 1188–94.
  19. Monbaliu E, La Peña D, Ortibus E, Molenaers G, Deklerck J, Feys H. Functional outcomes in children and young people with dyskinetic cerebral palsy. *Dev Med Child Neurol* 2017; **59**: 634–40.
  20. Narayanan UG, Fehlings D, Weir S, Knights S, Kiran S, Campbell K. Initial development and validation of the Caregiver Priorities and Child Health Index of Life with Disabilities (CPCCHILD). *Dev Med Child Neurol* 2006; **48**: 804–12.
  21. Novak I, Hines M, Goldsmith S, Barclay R. Clinical prognostic messages from a systematic review on cerebral palsy. *Pediatrics* 2012; **130**: e1285–312.
  22. Haberehner H, Bonouvié LA, Boeschoten K, et al. Use of the Dyskinesia Impairment Scale in non-ambulatory dyskinetic cerebral palsy. *Dev Med Child Neurol* 2020; **62**: 494–9.
  23. Stewart K, Harvey A. The Identification and Measurement of Dyskinesia in Children with Cerebral Palsy: A Toolkit for Clinicians. Melbourne: Murdoch Children's Research Institute, 2018.
  24. Kirshner B, Guyatt G. A methodological framework for assessing health indices. *J Chronic Diseases* 1985; **38**: 27–36.
  25. Monbaliu E, De Cock P, Mailleux L, Dan B, Feys H. The relationship of dystonia and choreoathetosis with activity, participation and quality of life in children and youth with dyskinetic cerebral palsy. *Eur J Paed Neurol* 2017; **21**: 327–35.
  26. Palisano R, Rosenbaum P, Walters SD, Russell D, Wood E, Galuppi B. Development and reliability of a system to classify gross motor function in children with cerebral palsy. *Dev Med Child Neurol* 1997; **39**: 214–23.
  27. Eliasson A-C, Kruminde-Sundholm L, Rösblad B, et al. The Manual Ability Classification System (MACS) for children with cerebral palsy: scale development and evidence of validity and reliability. *Dev Med Child Neurol* 2006; **48**: 549–54.
  28. Hidecker MJ, Paneth N, Rosenbaum PL, et al. Developing and validating the communication function classification system for individuals with cerebral palsy. *Dev Med Child Neurol* 2011; **53**: 704–10.
  29. Sellers D, Mandy A, Pennington L, Hankins M, Morris C. Development and reliability of a system to classify the eating and drinking ability of people with cerebral palsy. *Dev Med Child Neurol* 2014; **56**: 245–51.
  30. Love S, Gibson N, Blair E, Watson L. Cerebral palsy description form [Internet]. Sydney: Australian Cerebral Palsy Register, 2013. <https://cpregister.com/wp-content/uploads/2020/03/CP-Description-Form-Version-IV-21Feb2020.pdf> (accessed 21 February 2020).
  31. Hand C, Morley S, Adams M. Developing a questionnaire to measure patients' beliefs about inhaler treatment for asthma: tests of validity and reliability. *Primary Care Respir J* 2000; **9**: 12–5.
  32. Stewart K, Tavender E, Rice J, Harvey A. Identification, classification and assessment of dyskinesia in children with cerebral palsy: a survey of clinicians. *J Paed Child Health* 2018; **54**: 432–8.
  33. Keszei AP, Novak M, Streiner DL. Introduction to health measurement scales. *J Psychosomatic Res* 2010; **68**: 319–23.
  34. Rice J, Skuza P, Baker F, Russo R, Fehlings D. Identification and measurement of dystonia in cerebral palsy. *Dev Med Child Neurol* 2017; **59**: 1249–55.
  35. Gimeno H, Gordon A, Tustin K, Lin JP. Functional priorities in daily life for children and young people with dystonic movement disorders and their families. *Eur J Paed Neurol* 2013; **17**: 161–8.